arXiv:0704.2391v2 [math.AG] 14 Dec 2009

ORDINARY DIFFERENTIAL SYSTEMS IN DIMENSION THREE WITH AFFINE WEYL GROUP SYMMETRY OF TYPES $D_4^{(1)}, B_3^{(1)}, G_2^{(1)}, D_3^{(2)}$ AND $A_2^{(2)}$

YUSUKE SASANO

Abstract. We present a four-parameter family of ordinary differential systems in dimension three with affine Weyl group symmetry of type $D_4^{(1)}$. By obtaining its first integral, we can reduce this system to the second-order non-linear ordinary differential equations of Painlevé type. We also study this system restricted its parameters. Each system can be obtained by connecting some invariant divisors in the system of type $D_4^{(1)}$. Each system admits affine Weyl group symmetry of types $B_3^{(1)}, G_2^{(1)}, D_3^{(2)}$ and $A_2^{(2)}$, respectively. These symmetries, holomorphy conditions and invariant divisors are new.

1. Introduction

In this paper, we present a 4-parameter family of ordinary differential systems in dimension three with affine Weyl group symmetry of type $D_4^{(1)}$. By obtaining its first integral, we can reduce this system to the second-order non-linear ordinary differential equations of Painlevé type. This reduced system parametrizes the first-order ordinary differential equation:

$$\frac{dX}{dt} = \frac{b(t)}{2\eta} X(X+1)(X+1-\eta)(X-\eta), \quad b(t) \in \mathbb{C}(t),\ \eta \in \mathbb{C} - \{0\}. \tag{1}$$

We also study this system restricted its parameters. Each system admits affine Weyl group symmetry of types $B_3^{(1)}, G_2^{(1)}, D_3^{(2)}$ and $A_2^{(2)}$, respectively.

Each system can be obtained by connecting some invariant divisors in the system of type $D_4^{(1)}$.

The Bäcklund transformations of each system satisfy

$$s_i(g) = g + \frac{\alpha_i}{f_i}\{f_i, g\} + \frac{1}{2!}\left(\frac{\alpha_i}{f_i}\right)^2 \{f_i, \{f_i, g\}\} + \cdots \quad (g \in \mathbb{C}(t)[x,y,z]), \tag{2}$$

where poisson bracket $\{,\}$ satisfies the relations:

$$\{z, x\} = \{z, y\} = 1, \quad \{x, y\} = 0. \tag{3}$$

Since these Bäcklund transformations have Lie theoretic origin, similarity reduction of a Drinfeld-Sokolov hierarchy admits such a Bäcklund symmetry.

These symmetries, holomorphy conditions and invariant divisors are new.

2000 *Mathematics Subjet Classification.* 34M55; 34M45; 58F05; 32S65.

Key words and phrases. Birational symmetry, Chazy equations, Painlevé equations.

2. THE SYSTEM OF TYPE $D_4^{(1)}$

In this paper, we study the third-order ordinary differential system:

$$
\left\{
\begin{aligned}
\frac{2\eta}{b(t)}\frac{dx}{dt} =& -2x(y-1)yz(x-\eta)(2x-2y+1-\eta)-2(2\alpha_0+2\alpha_1+4\alpha_2+\alpha_3+\alpha_4)x^3y\\
&-2(\alpha_0+\alpha_1)xy^3+(5\alpha_0+5\alpha_1+6\alpha_2+\alpha_3+\alpha_4)x^2y^2\\
&-\{5\alpha_0+5\alpha_1+6\alpha_2+2\alpha_3-3(2\alpha_0+2\alpha_1+4\alpha_2+\alpha_3+\alpha_4)\eta\}x^2y\\
&+\{3(\alpha_0+\alpha_1)-(4\alpha_0+6\alpha_1+6\alpha_2+\alpha_3+\alpha_4)\eta\}xy^2+x^4\\
&+2(\alpha_0+\alpha_1+2\alpha_2+\alpha_3-\eta)x^3+2\alpha_1\eta y^3\\
&+\{(\alpha_0+\alpha_1+2\alpha_2+\alpha_3)(\eta^2-3\eta+1)+\alpha_4\eta^2\}x^2+\alpha_1(\eta-3)\eta y^2\\
&+\{-(\alpha_0+\alpha_1)+2(2\alpha_0+3\alpha_1+3\alpha_2+\alpha_3)\eta-(2\alpha_0+2\alpha_1+4\alpha_2+\alpha_3+\alpha_4)\eta^2\}xy\\
&+(\alpha_0+\alpha_1+2\alpha_2+\alpha_3)(\eta-1)\eta x-\alpha_1(\eta-1)\eta y,\\
-\frac{2\eta}{b(t)}\frac{dy}{dt} =& 2x(y-1)yz(x-\eta)(2x-2y+1-\eta)-2(\alpha_3+\alpha_4)x^3y\\
&-2xy^3(\alpha_0+\alpha_1+4\alpha_2+2\alpha_3+2\alpha_4)+x^2y^2(\alpha_0+\alpha_1+6\alpha_2+5\alpha_3+5\alpha_4)\\
&+x^2y\{-(\alpha_0+\alpha_1+6\alpha_2+4\alpha_3+6\alpha_4)+3(\alpha_3+\alpha_4)\eta\}\\
&+xy^2\{3(\alpha_0+\alpha_1+4\alpha_2+2\alpha_3+2\alpha_4)-(2\alpha_0+6\alpha_2+5\alpha_3+5\alpha_4)\eta\}+y^4+2\alpha_4x^3\\
&+2y^3((\alpha_0+2\alpha_2+\alpha_3+\alpha_4)\eta-1)-\alpha_4(3\eta-1)x^2\\
&+y^2\{1-3(\alpha_0+2\alpha_2+\alpha_3+\alpha_4)\eta+(\alpha_0+2\alpha_2+\alpha_3+\alpha_4)\eta^2\}\\
&+xy\{-(\alpha_0+\alpha_1+4\alpha_2+2\alpha_3+2\alpha_4)+2(\alpha_0+3\alpha_2+2\alpha_3+3\alpha_4)\eta-(\alpha_3+\alpha_4)\eta^2\}\\
&+\alpha_4(\eta-1)\eta x-(\alpha_0+2\alpha_2+\alpha_3+\alpha_4)(\eta-1)\eta y,\\
\frac{2\eta}{b(t)}\frac{dz}{dt} =& (2x-2y+1-\eta)\{2x^2y+2xy^2-x^2-y^2\eta-2(\eta+1)xy+\eta(x+y)\}z^2\\
&+z[-2x^3(\alpha_3+\alpha_4)+2y^3(\alpha_0+\alpha_1)+2x^2y(\alpha_0+\alpha_1+6\alpha_2+2\alpha_3+2\alpha_4)\\
&-2xy^2(2\alpha_0+2\alpha_1+6\alpha_2+\alpha_3+\alpha_4)+x^2\{-(\alpha_0+\alpha_1+6\alpha_2+4\alpha_3)+3(\alpha_3+\alpha_4)\eta\}\\
&+y^2(-3(\alpha_0+\alpha_1)+(4\alpha_0+6\alpha_2+\alpha_3+\alpha_4)\eta)\\
&-4xy\{-(\alpha_0+\alpha_1+3\alpha_2+\alpha_3)+(\alpha_0+3\alpha_2+\alpha_3+\alpha_4)\eta\}\\
&+x\{-(\alpha_0+\alpha_1+4\alpha_2+2\alpha_3)+2(\alpha_0+3\alpha_2+2\alpha_3)\eta-(\alpha_3+\alpha_4)\eta^2\}\\
&+y\{\alpha_0+\alpha_1-2(2\alpha_0+3\alpha_2+\alpha_3)\eta+(2\alpha_0+4\alpha_2+\alpha_3+\alpha_4)\eta^2\}\\
&-(\alpha_0+2\alpha_2+\alpha_3)(\eta-1)\eta]+\alpha_2[(\alpha_0+\alpha_1+2\alpha_2-\alpha_3-\alpha_4)x^2\\
&+(\alpha_0+\alpha_1-2\alpha_2-\alpha_3-\alpha_4)y^2-2(\alpha_0+\alpha_1-\alpha_3-\alpha_4)xy\\
&+x\{\alpha_0+\alpha_1-2\alpha_3-(2\alpha_0+2\alpha_2-\alpha_3-\alpha_4)\eta\}\\
&+y\{-\alpha_0-\alpha_1+2\alpha_2+2\alpha_3+(2\alpha_0-\alpha_3-\alpha_4)\eta\}+(\eta-1)(\alpha_2+\alpha_3+(\alpha_0+\alpha_2)\eta)].
\end{aligned}
\right.
\tag{4}
$$

Here x, y and z denote unknown complex variables, $b(t) \in \mathbb{C}(t)$, $\eta \in \mathbb{C} - \{0\}$, and $\alpha_0, \alpha_1, \dots, \alpha_4$ are complex parameters satisfying the relation:

$$\alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4 = 1. \tag{5}$$

Theorem 2.1. *Let us consider the following ordinary differential system in the polynomial class*:

$$\begin{cases} \dfrac{dx}{dt} = f_1(x, y, z), \\ \dfrac{dy}{dt} = f_2(x, y, z), \\ \dfrac{dz}{dt} = f_3(x, y, z). \end{cases}$$

We assume that

$(A1)$ $deg(f_i) = 6$ with respect to x, y, z.

$(A2)$ The right-hand side of this system becomes again a polynomial in each coordinate system (x_i, y_i, z_i) $(i = 0, 1, \dots, 4)$.

$$\begin{aligned} &0)\ x_0 = x - \eta, \quad y_0 = y, \quad z_0 = z - \frac{\alpha_0}{x - \eta}, \\ &1)\ x_1 = x, \quad y_1 = y, \quad z_1 = z - \frac{\alpha_1}{x}, \\ &2)\ x_2 = x + \frac{\alpha_2}{z}, \quad y_2 = y + \frac{\alpha_2}{z}, \quad z_2 = z, \\ &3)\ x_3 = x, \quad y_3 = y - 1, \quad z_3 = z - \frac{\alpha_3}{y - 1}, \\ &4)\ x_4 = x, \quad y_4 = y, \quad z_4 = z - \frac{\alpha_4}{y}. \end{aligned} \tag{6}$$

Then such a system coincides with the system (4).

These transition functions satisfy the condition:

$$dx_i \wedge dy_i \wedge dz_i = dx \wedge dy \wedge dz \quad (i = 0, 1, \dots, 4).$$

Theorem 2.2. *The system* (4) *admits the affine Weyl group symmetry of type* $D_4^{(1)}$ *as the group of its Bäcklund transformations, whose generators are explicitly given as follows*:

with the notation $(*) := (x, y, z; \alpha_0, \alpha_1, \dots, \alpha_4)$,

$$\begin{aligned}
s_0 &: (*) \to \left(x, y, z - \frac{\alpha_0}{x-\eta}; -\alpha_0, \alpha_1, \alpha_2 + \alpha_0, \alpha_3, \alpha_4\right),\\
s_1 &: (*) \to \left(x, y, z - \frac{\alpha_1}{x}; \alpha_0, -\alpha_1, \alpha_2 + \alpha_1, \alpha_3, \alpha_4\right),\\
s_2 &: (*) \to \left(x + \frac{\alpha_2}{z}, y + \frac{\alpha_2}{z}, z; \alpha_0 + \alpha_2, \alpha_1 + \alpha_2, -\alpha_2, \alpha_3 + \alpha_2, \alpha_4 + \alpha_2\right),\\
s_3 &: (*) \to \left(x, y, z - \frac{\alpha_3}{y-1}; \alpha_0, \alpha_1, \alpha_2 + \alpha_3, -\alpha_3, \alpha_4\right),\\
s_4 &: (*) \to \left(x, y, z - \frac{\alpha_4}{y}; \alpha_0, \alpha_1, \alpha_2 + \alpha_4, \alpha_3, -\alpha_4\right).
\end{aligned}$$

We note that the generators $s_0, s_1, \dots, s_4$ are determined by the invariant divisors (2.3) (see next proposition).

PROPOSITION 2.3. *The system* (4) *has the following invariant divisors*:

parameter's relation	f_i
$\alpha_0 = 0$	$f_0 := x - \eta$
$\alpha_1 = 0$	$f_1 := x$
$\alpha_2 = 0$	$f_2 := z$
$\alpha_3 = 0$	$f_3 := y - 1$
$\alpha_4 = 0$	$f_4 := y$

We note that when $\alpha_1 = 0$, we see that the system (4) admits a particular solution $x = 0$.

3. REDUCTION OF THE $D_4^{(1)}$ SYSTEM

In this section, we show that the system (4) has its first integral. Thanks to this first integral, we can reduce the system (4) to the second-order non-linear ordinary differential equations.

PROPOSITION 3.1. *The system* (4) *has its first integral*:

$$\frac{d(x-y)}{dt} = \frac{b(t)}{2\eta}(x-y)(x-y+1)(x-y+1-\eta)(x-y-\eta). \tag{7}$$

Under the condition

$$b(t) = \frac{2\eta}{t(t-1)(t+\eta)(t+\eta-1)}, \tag{8}$$

the equation (7) admits a particular solution:

$$x = y - t. \tag{9}$$

THEOREM 3.2. *Under the conditions*

$$b(t)=\frac{2\eta}{t(t-1)\{t^2+(2\eta-1)t+\eta(\eta-1)\}}, \tag{10}$$

$$x=y-t, \tag{11}$$

for the system (4) *we make the change of parameters and variables*

$$\alpha_0=A_1, \quad \alpha_1=A_0, \quad \alpha_2=A_2, \quad \alpha_3=A_3, \quad \alpha_4=A_4, \quad X:=y, \quad Y:=z \tag{12}$$

from $\alpha_0,\alpha_1,\dots,\alpha_4,x,y,z$ *to* A_i,X,Y. *Then the system* (4) *can also be written in the new variables* X,Y *and parameters* A_i *as a Hamiltonian system. This new system tends to*

$$\frac{dX}{dt}=\frac{\partial H_{VI}}{\partial Y}, \quad \frac{dY}{dt}=-\frac{\partial H_{VI}}{\partial X} \tag{13}$$

with the polynomial Hamiltonian

$$\begin{aligned}&H_{VI}(X,Y,t;A_1,A_0,A_2,A_3,A_4)\\ &=\frac{1}{t(t-1)}[Y^2(X-t)(X-1)X-\{(A_1-1)(X-1)X+A_3(X-t)X\\ &+A_4(X-t)(X-1)\}Y+A_2(A_0+A_2)X]\end{aligned} \tag{14}$$

as $\eta\to\infty$.

This system is the Painlevé VI system.

We also see that the equation (7) admits a particular solution:

$$x=y. \tag{15}$$

THEOREM 3.3. *Under the condition*

$$x=y, \tag{16}$$

for the system (4) *we make the change of variables*

$$X:=y, \quad Y:=z \tag{17}$$

from x,y,z *to* X,Y. *Then the system* (4) *can also be written in the new variables* X,Y *as a Hamiltonian system. This new system tends to*

$$\frac{dX}{dt}=\frac{\partial H}{\partial Y}, \quad \frac{dY}{dt}=-\frac{\partial H}{\partial X} \tag{18}$$

with the polynomial Hamiltonian

(19)

$$\begin{aligned}&H(X,Y,t;\alpha_0,\alpha_1,\alpha_2,\alpha_3,\alpha_4)\\ &=-\frac{\eta-1}{2\eta}b(t)[-(X-1)(X-\eta)X^2Y^2-\{2\alpha_2(X-1)(X-\eta)-\alpha_0\eta(X-1)-\alpha_3(X-\eta)\}XY\\ &-{\alpha_2}^2X^2+\alpha_2\{1-(\alpha_0+\alpha_1+\alpha_2+\alpha_4)+(1-(\alpha_1+\alpha_2+\alpha_3+\alpha_4))\eta\}X].\end{aligned}$$

Elimination of Y from the system (18) gives the second-order non-linear ordinary differential equation for the variable X:

$$\begin{aligned}\frac{d^2X}{dt^2} &= \left\{\frac{1}{2(X-1)}+\frac{1}{X}+\frac{1}{2(X-\eta)}\right\}\left(\frac{dX}{dt}\right)^2+\frac{\frac{db(t)}{dt}}{b(t)}\frac{dX}{dt}\\ &\quad-\frac{b(t)^2}{8\eta^2(X-1)(X-\eta)}[(\eta-1)^3X^2\{(\alpha_0\eta+\alpha_3)X-(1-(\alpha_1+2\alpha_2+\alpha_4))\eta\}\\ &\quad\times\{(\alpha_0\eta-\alpha_3)X+(1-(2\alpha_0+\alpha_1+2\alpha_2+\alpha_4))\eta\}].\end{aligned} \tag{20}$$

Proposition 3.4. *The system* (18) *admits the affine Weyl group symmetry of type* $D_4^{(1)}$ *as the group of its Bäcklund transformations, whose generators are explicitly given as follows*: *with the notation* $(*) := (X, Y; \alpha_0, \alpha_1, \dots, \alpha_4)$,

$$\begin{aligned}
s_0 &: (*) \rightarrow \left(X, Y-\frac{\alpha_0}{X-\eta}; -\alpha_0, \alpha_1, \alpha_2+\alpha_0, \alpha_3, \alpha_4\right),\\
s_1 &: (*) \rightarrow \left(X, Y-\frac{\alpha_1}{X}; \alpha_0, -\alpha_1, \alpha_2+\alpha_1, \alpha_3, \alpha_4\right),\\
s_2 &: (*) \rightarrow \left(X+\frac{\alpha_2}{Y}, Y; \alpha_0+\alpha_2, \alpha_1+\alpha_2, -\alpha_2, \alpha_3+\alpha_2, \alpha_4+\alpha_2\right),\\
s_3 &: (*) \rightarrow \left(X, Y-\frac{\alpha_3}{X-1}; \alpha_0, \alpha_1, \alpha_2+\alpha_3, -\alpha_3, \alpha_4\right),\\
s_4 &: (*) \rightarrow \left(X, Y-\frac{\alpha_4}{X}; \alpha_0, \alpha_1, \alpha_2+\alpha_4, \alpha_3, -\alpha_4\right).
\end{aligned}$$

4. The system of type $B_3^{(1)}$

In this section, we present a 3-parameter family of ordinary differential systems in dimension three with affine Weyl group symmetry of type $B_3^{(1)}$. This system is equivalent to the system (4) restricted its parameters.

This system can be obtained by connecting the invariant divisors x and y in the system (4).

For this system, we can discuss some reductions to the second-order ordinary differential equations in the same way in previous section because this system is equivalent to the system (4) only restricted its parameters.

Theorem 4.1. *Let us consider the following ordinary differential system in the polynomial class*:

$$\begin{cases}\dfrac{dx}{dt}=f_1(x,y,z),\\ \dfrac{dy}{dt}=f_2(x,y,z),\\ \dfrac{dz}{dt}=f_3(x,y,z).\end{cases}$$

We assume that

$(A1)$ $deg(f_i)=6$ with respect to x, y, z.

$(A2)$ The right-hand side of this system becomes again a polynomial in each coordinate system (x_i, y_i, z_i) $(i = 0, 1, 2, 3)$.

$$
\begin{aligned}
&0)\ x_0 = x - \eta, \quad y_0 = y, \quad z_0 = z - \frac{\beta_0}{x - \eta},\\
&1)\ x_1 = x, \quad y_1 = y - 1, \quad z_1 = z - \frac{\beta_1}{y - 1},\\
&2)\ x_2 = x + \frac{\beta_2}{z}, \quad y_2 = y + \frac{\beta_2}{z}, \quad z_2 = z,\\
&3)\ x_3 = x, \quad y_3 = y, \quad z_3 = z - \frac{\beta_3(x + y)}{xy}.
\end{aligned}
\tag{21}
$$

Then such a system coincides with the system (4) with the parameter's relations:

$$\alpha_4 = \alpha_1, \quad \beta_0 := \alpha_0, \quad \beta_1 := \alpha_3, \quad \beta_2 := \alpha_2, \quad \beta_3 := \alpha_1. \tag{22}$$

Here, the complex parameters $\beta_0, \beta_1, \beta_2, \beta_3$ satisfy the relation:

$$\beta_0 + \beta_1 + 2\beta_2 + 2\beta_3 = 1. \tag{23}$$

These transition functions satisfy the condition:

$$dx_i \wedge dy_i \wedge dz_i = dx \wedge dy \wedge dz \quad (i = 0, 1, 2, 3).$$

Theorem 4.2. *This system admits the affine Weyl group symmetry of type $B_3^{(1)}$ as the group of its Bäcklund transformations, whose generators are explicitly given as follows: with the notation $(*) := (x, y, z; \beta_0, \beta_1, \beta_2, \beta_3)$,*

$$
\begin{aligned}
&s_0 : (*) \rightarrow \left(x, y, z - \frac{\beta_0}{x - \eta}; -\beta_0, \beta_1, \beta_2 + \beta_0, \beta_3\right),\\
&s_1 : (*) \rightarrow \left(x, y, z - \frac{\beta_1}{y - 1}; \beta_0, -\beta_1, \beta_2 + \beta_1, \beta_3\right),\\
&s_2 : (*) \rightarrow \left(x + \frac{\beta_2}{z}, y + \frac{\beta_2}{z}, z; \beta_0 + \beta_2, \beta_1 + \beta_2, -\beta_2, \beta_3 + \beta_2\right),\\
&s_3 : (*) \rightarrow \left(x, y, z - \frac{\beta_3(x + y)}{xy}; \beta_0, \beta_1, \beta_2 + 2\beta_3, -\beta_3\right).
\end{aligned}
$$

The Bäcklund transformations of each system satisfy

$$s_i(g) = g + \frac{\alpha_i}{f_i}\{f_i, g\} + \frac{1}{2!}\left(\frac{\alpha_i}{f_i}\right)^2 \{f_i, \{f_i, g\}\} + \cdots \quad (g \in \mathbb{C}(t)[x, y, z]), \tag{24}$$

where poisson bracket $\{,\}$ satisfies the relations:

$$\{z, x\} = \{z, y\} = 1, \quad \{x, y\} = 0. \tag{25}$$

Since these Bäcklund transformations have Lie theoretic origin, similarity reduction of a Drinfeld-Sokolov hierarchy admits such a Bäcklund symmetry.

We note that all generators s_0, s_1, s_2, s_3 are determined by the invariant divisors (4.3) (see next proposition).

PROPOSITION 4.3. *This system has the following invariant divisors*:

parameter's relation	f_i
$\beta_0=0$	$f_0:=x-\eta$
$\beta_1=0$	$f_1:=y-1$
$\beta_2=0$	$f_2:=z$
$\beta_3=0$	$f_3:=xy$

We note that when $\beta_3=0$, after we make the birational transformations:

$$x_3=xy,\ y_3=y,\ z_3=z \tag{26}$$

we see that in the coordinate system (x_3,y_3,z_3) the system admits a particular solution $x_3=0$.

5. THE SYSTEM OF TYPE $D_3^{(2)}$

In this section, we present a 2-parameter family of ordinary differential systems in dimension three with affine Weyl group symmetry of type $D_3^{(2)}$. This system is equivalent to the system (4) restricted its parameters.

THEOREM 5.1. *Let us consider the following ordinary differential system in the polynomial class*:

$$\begin{cases} \dfrac{dx}{dt}=f_1(x,y,z),\\ \dfrac{dy}{dt}=f_2(x,y,z),\\ \dfrac{dz}{dt}=f_3(x,y,z). \end{cases}$$

We assume that

$(A1)$ $deg(f_i)=6$ with respect to x,y,z.

$(A2)$ The right-hand side of this system becomes again a polynomial in each coordinate system (x_i,y_i,z_i) $(i=1,2,3)$.

$$\begin{aligned} &1)\ x_1=x-\eta,\quad y_1=y-1,\quad z_1=z-\frac{\beta_0(x+y-\eta-1)}{(x-\eta)(y-1)},\\ &2)\ x_2=x+\frac{\beta_1}{z},\quad y_2=y+\frac{\beta_1}{z},\quad z_2=z,\\ &3)\ x_3=x,\quad y_3=y,\quad z_3=z-\frac{\beta_2(x+y)}{xy}. \end{aligned} \tag{27}$$

Then such a system coincides with the system (4) with the parameter's relations:

$$\alpha_4=\alpha_1,\quad \alpha_0=\alpha_3,\quad \beta_0:=\alpha_1,\quad \beta_1:=\alpha_2,\quad \beta_2:=\alpha_3. \tag{28}$$

Here, the complex parameters β_0,β_1,β_2 satisfy the relation:

$$\beta_0+\beta_1+\beta_2=\frac{1}{2}. \tag{29}$$

These transition functions satisfy the condition:

$$dx_i \wedge dy_i \wedge dz_i = dx \wedge dy \wedge dz \quad (i = 1, 2, 3).$$

THEOREM 5.2. *This system admits the affine Weyl group symmetry of type $D_3^{(2)}$ as the group of its Bäcklund transformations, whose generators are explicitly given as follows: with the notation $(*) := (x, y, z; \beta_0, \beta_1, \beta_2)$,*

$$\begin{aligned}
s_0 : (*) &\rightarrow \left(x, y, z - \frac{\beta_0(x + y - \eta - 1)}{(x - \eta)(y - 1)}; -\beta_0, \beta_1 + 2\beta_0, \beta_2\right),\\
s_1 : (*) &\rightarrow \left(x + \frac{\beta_1}{z}, y + \frac{\beta_1}{z}, z; \beta_0 + \beta_1, -\beta_1, \beta_2 + \beta_1\right),\\
s_2 : (*) &\rightarrow \left(x, y, z - \frac{\beta_2(x + y)}{xy}; \beta_0, \beta_1 + 2\beta_2, -\beta_2\right).
\end{aligned}$$

We note that the generators s_0, s_1, s_2 are determined by the invariant divisors (5.3) (see next proposition).

PROPOSITION 5.3. *This system has the following invariant divisors:*

parameter's relation	f_i
$\beta_0 = 0$	$f_0 := (x - \eta)(y - 1)$
$\beta_1 = 0$	$f_1 := z$
$\beta_2 = 0$	$f_2 := xy$

6. THE SYSTEM OF TYPE $G_2^{(1)}$

In this section, we present a 2-parameter family of ordinary differential systems in dimension three with affine Weyl group symmetry of type $G_2^{(1)}$. This system is equivalent to the system (4) restricted its parameters

$$\alpha_4 = \alpha_3 = \alpha_1, \quad \beta_0 := \alpha_0, \quad \beta_1 := \alpha_2, \quad \beta_2 := \alpha_3. \tag{30}$$

Here, the complex parameters $\beta_0, \beta_1, \beta_2$ satisfy the relation:

$$\beta_0 + 2\beta_1 + 3\beta_2 = 1. \tag{31}$$

THEOREM 6.1. *This system admits the affine Weyl group symmetry of type $G_2^{(1)}$ as the group of its Bäcklund transformations, whose generators are explicitly given as follows: with the notation $(*) := (x, y, z; \beta_0, \beta_1, \beta_2)$,*

$$\begin{aligned}
s_0 : (*) &\rightarrow \left(x, y, z - \frac{\beta_0}{x - \eta}; -\beta_0, \beta_1 + \beta_0, \beta_2\right),\\
s_1 : (*) &\rightarrow \left(x + \frac{\beta_1}{z}, y + \frac{\beta_1}{z}, z; \beta_0 + \beta_1, -\beta_1, \beta_2 + \beta_1\right),\\
s_2 : (*) &\rightarrow \left(x, y, z - \frac{\beta_2\{y(y - 1) + x(y - 1) + xy\}}{xy(y - 1)}; \beta_0, \beta_1 + 3\beta_2, -\beta_2\right).
\end{aligned}$$

We note that the generators s_0, s_1, s_2 are determined by the invariant divisors (6.2) (see next proposition).

PROPOSITION 6.2. *This system has the following invariant divisors*:

parameter's relation	f_i
$\beta_0 = 0$	$f_0 := x - \eta$
$\beta_1 = 0$	$f_1 := z$
$\beta_2 = 0$	$f_2 := xy(y-1)$

7. THE SYSTEM OF TYPE $A_2^{(2)}$

In this section, we present a 1-parameter family of ordinary differential systems in dimension three with affine Weyl group symmetry of type $A_2^{(2)}$. This system is equivalent to the system (4) restricted its parameters

$$\alpha_4 = \alpha_3 = \alpha_0 = \alpha_1, \quad \beta_0 := \alpha_2, \quad \beta_1 := \alpha_1. \tag{32}$$

Here, the complex parameters β_0, β_1 satisfy the relation:

$$\beta_0 + 2\beta_1 = \frac{1}{2}. \tag{33}$$

THEOREM 7.1. *This system admits the affine Weyl group symmetry of type $A_2^{(2)}$ as the group of its Bäcklund transformations, whose generators are explicitly given as follows*: *with the notation* $(*) := (x, y, z; \beta_0, \beta_1)$,

$$\begin{aligned} s_0 : (*) &\rightarrow \left(x + \frac{\beta_0}{z}, y + \frac{\beta_0}{z}, z; -\beta_0, \beta_1 + \beta_0\right), \\ s_1 : (*) &\rightarrow \Bigl(x, y, z - \frac{\beta_1\{y(x-\eta)(y-1) + x(x-\eta)(y-1) + xy(y-1) + xy(x-\eta)\}}{xy(x-\eta)(y-1)}; \\ &\qquad \beta_0 + 4\beta_1, -\beta_1\Bigr). \end{aligned}$$

We note that the generators s_0, s_1 are determined by the invariant divisors (7.2) (see next proposition).

PROPOSITION 7.2. *This system has the following invariant divisors*:

parameter's relation	f_i
$\beta_0 = 0$	$f_0 := z$
$\beta_1 = 0$	$f_1 := xy(x-\eta)(y-1)$

8. APPENDIX A

It is well-known that the fifth Painlevé equation has symmetries under the affine Weyl group of type $A_3^{(1)}$. In this section, we present a 3-parameter family of the systems of the first-order ordinary differential equations.

THEOREM 8.1. *The fifth Painlevé equation is equivalent to a 3-parameter family of the systems of the first-order ordinary differential equations*:

$$\left\{\begin{aligned}\frac{df_0}{dt}&=-\frac{\varphi}{2}(-2f_0f_1f_2+af_0f_2+(\alpha_0+\alpha_1+\alpha_3)f_0-\alpha_0f_2),\\ \frac{df_1}{dt}&=\frac{\varphi}{2}(-f_0f_1^2-f_1^2f_2+af_0f_1+af_1f_2-a\alpha_1+(\alpha_1+\alpha_3)f_1),\\ \frac{df_2}{dt}&=-\frac{\varphi}{2}(-2f_0f_1f_2+af_0f_2+(\alpha_1+\alpha_2+\alpha_3)f_2-\alpha_2f_0).\end{aligned}\right. \tag{34}$$

Here, f_0, f_1 and f_2 denote unknown complex variables and $\alpha_0, \alpha_1, \alpha_2, \alpha_3$ and a are constant complex parameters with $\alpha_0+\alpha_1+\alpha_2+\alpha_3=1$ and φ is a nonzero parameter which can be fixed arbitrarily.

We see that the system (34) has its first integral:

$$\frac{d(f_2-f_0)}{dt}=f_2-f_0. \tag{35}$$

We can solve this equation by

$$f_2-f_0=e^{(t+c)}. \tag{36}$$

Here we set

$$t+c=logT,\ x:=f_0,\ y:=f_1, \tag{37}$$

then we can obtain the fifth Painlevé system:

$$\left\{\begin{aligned}\frac{dx}{dT}&=\frac{\partial H_V}{\partial y}=-\frac{2x^2y}{T}+\frac{ax^2}{T}-2xy+\left(a+\frac{\alpha_1+\alpha_3}{T}\right)x-\alpha_0,\\ \frac{dy}{dT}&=-\frac{\partial H_V}{\partial x}=\frac{2xy^2}{T}+y^2-\frac{2axy}{T}-\left(a+\frac{\alpha_1+\alpha_3}{T}\right)y+\frac{a\alpha_1}{T}\end{aligned}\right. \tag{38}$$

with the polynomial Hamiltonian

$$H_V=-\frac{x^2y^2}{T}+\frac{ax^2y}{T}-xy^2+\left(a+\frac{\alpha_1+\alpha_3}{T}\right)xy-\alpha_0y-\frac{a\alpha_1}{T}x. \tag{39}$$

THEOREM 8.2. *This system admits the affine Weyl group symmetry of type $A_3^{(1)}$ as the group of its Bäcklund transformations, whose generators s_0, s_1, s_2, s_3, π are explicitly given* *as follows*:

(40)

$$\begin{aligned} s_0 :&(f_0,f_1,f_2;\alpha_0,\alpha_1,\alpha_2,\alpha_3) \rightarrow \left(f_0, f_1+\frac{\alpha_0}{f_0}, f_2; -\alpha_0, \alpha_1+\alpha_0, \alpha_2, \alpha_3+\alpha_0\right),\\ s_1 :&(f_0,f_1,f_2;\alpha_0,\alpha_1,\alpha_2,\alpha_3) \rightarrow \left(f_0-\frac{\alpha_1}{f_1}, f_1, f_2-\frac{\alpha_1}{f_1}; \alpha_0+\alpha_1, -\alpha_1, \alpha_2+\alpha_1, \alpha_3\right),\\ s_2 :&(f_0,f_1,f_2;\alpha_0,\alpha_1,\alpha_2,\alpha_3) \rightarrow \left(f_0, f_1+\frac{\alpha_2}{f_2}, f_2; \alpha_0, \alpha_1+\alpha_2, -\alpha_2, \alpha_3+\alpha_2\right),\\ s_3 :&(f_0,f_1,f_2;\alpha_0,\alpha_1,\alpha_2,\alpha_3) \rightarrow \left(f_0-\frac{\alpha_3}{f_1-a}, f_1, f_2-\frac{\alpha_3}{f_1-a}; \alpha_0+\alpha_3, \alpha_1, \alpha_2+\alpha_3, -\alpha_3\right),\\ \pi :&(f_0,f_1,f_2;\alpha_0,\alpha_1,\alpha_2,\alpha_3) \rightarrow (f_2,f_1,f_0;\eta,\alpha_2,\alpha_1,\alpha_0,\alpha_3). \end{aligned}$$

The Bäcklund transformations of each system satisfy

$$s_i(g) = g + \frac{\alpha_i}{f_i}\{f_i,g\} + \frac{1}{2!}\left(\frac{\alpha_i}{f_i}\right)^2\{f_i,\{f_i,g\}\} + \cdots \quad (g \in \mathbb{C}[f_0,f_1,f_2]), \tag{41}$$

where poisson bracket $\{,\}$ satisfies the relations:

$$\{f_0,f_1\} = \{f_2,f_1\} = 1, \quad \{f_0,f_2\} = 0. \tag{42}$$

Since these Bäcklund transformations have Lie theoretic origin, similarity reduction of a Drinfeld-Sokolov hierarchy admits such a Bäcklund symmetry.

9. APPENDIX B

It is well-known that the third Painlevé equation has symmetries under the affine Weyl group of type $C_2^{(1)}$. In this section, we present a new representation of the third Painlevé equation.

THEOREM 9.1. *The third Painlevé equation can be written in the following symmetric form*:

$$\begin{cases} \dfrac{df_0}{dt} = -2f_0f_1f_2 + (\alpha_0+2\alpha_1)f_0 - \alpha_0 f_2,\\ \dfrac{df_1}{dt} = (f_0+f_2)f_1^2 - 2\alpha_1 f_1 + \eta,\\ \dfrac{df_2}{dt} = -2f_0f_1f_2 + (2\alpha_1+\alpha_2)f_2 - \alpha_2 f_0. \end{cases} \tag{43}$$

Here, f_0, f_1 and f_2 denote unknown complex variables and α_0, α_1 and α_2 are constant parameters with $\alpha_0 + 2\alpha_1 + \alpha_2 = 1$ and η is a nonzero parameter which can be fixed arbitrarily.

We see that the system (43) has its first integral:

$$\frac{d(f_0-f_2)}{dt} = f_0 - f_2. \tag{44}$$

We can solve this equation by

$$f_0 - f_2 = e^{(t+c)}. \tag{45}$$

Here we set

$$t + c = logT, \ x := \frac{1}{f_1}, \ y := -(f_1 f_2 + \alpha_2) f_1, \tag{46}$$

then we can obtain the third Painlevé system:

$$\begin{cases} \dfrac{dx}{dT} = \dfrac{\partial H_{III}}{\partial y} = \dfrac{2x^2y}{T} - \dfrac{\eta x^2}{T} + \dfrac{2(\alpha_1+\alpha_2)x}{T} - 1, \\ \dfrac{dy}{dT} = -\dfrac{\partial H_{III}}{\partial x} = -\dfrac{2xy^2}{T} + \dfrac{2\eta xy}{T} - \dfrac{2(\alpha_1+\alpha_2)y}{T} + \dfrac{\eta\alpha_2}{T} \end{cases} \tag{47}$$

with the polynomial Hamiltonian

$$H_{III} = \frac{x^2y^2 - \eta x^2 y + 2(\alpha_1+\alpha_2)xy - Ty - \eta\alpha_2 x}{T}. \tag{48}$$

THEOREM 9.2. *This system admits extended affine Weyl group symmetry of type* $C_2^{(1)}$ *as the group of its Bäcklund transformations, whose generators* s_0, s_1, s_2, π *are explicitly given as follows*:

$$\begin{aligned} s_0 : (f_0, f_1, f_2; \eta, \alpha_0, \alpha_1, \alpha_2) \rightarrow & \left(f_0, f_1 + \frac{\alpha_0}{f_0}, f_2; \eta, -\alpha_0, \alpha_1 + \alpha_0, \alpha_2\right), \\ s_1 : (f_0, f_1, f_2; \eta, \alpha_0, \alpha_1, \alpha_2) \rightarrow & (f_0 - \frac{2\alpha_1}{f_1} + \frac{\eta}{f_1^2}, f_1, f_2 - \frac{2\alpha_1}{f_1} + \frac{\eta}{f_1^2}; \\ & \qquad - \eta, \alpha_0 + 2\alpha_1, -\alpha_1, \alpha_2 + 2\alpha_1), \\ s_2 : (f_0, f_1, f_2; \eta, \alpha_0, \alpha_1, \alpha_2) \rightarrow & \left(f_0, f_1 + \frac{\alpha_2}{f_2}, f_2; \eta, \alpha_0, \alpha_1 + \alpha_2, -\alpha_2\right), \\ \pi : (f_0, f_1, f_2; \eta, \alpha_0, \alpha_1, \alpha_2) \rightarrow & (f_2, f_1, f_0; \eta, \alpha_2, \alpha_1, \alpha_0). \end{aligned} \tag{49}$$



FIGURE 1. The transformations described in Theorem 9.2 satisfy the relations: ${s_0}^2 = {s_1}^2 = {s_2}^2 = \pi^2 = (s_0 s_2)^2 = (s_0 s_1)^4 = (s_1 s_2)^4 = 1, \ \pi(s_0, s_1, s_2) = (s_2, s_1, s_0)\pi$.

Theorem 9.3. *For the system* (34) *of type* $A_3^{(1)}$*, we make the change of parameters*

$$\alpha_0=\beta_0,\ \alpha_1=-\frac{\eta}{a},\ \alpha_2=\beta_2,\ \alpha_3=2\beta_1+\frac{\eta}{a},\ \varphi=-2, \tag{50}$$

from $\alpha_0,\alpha_1,\alpha_2,\alpha_3$ *to* β_0,β_1,β_2*. This new system tends to the system* (43) *of type* $C_2^{(1)}$ *as* $a\to 0$.

We note that

$$\alpha_0+\alpha_1+\alpha_2+\alpha_3=\beta_0+2\beta_1+\beta_2=1.$$

References

[1] P. Painlevé, *Mémoire sur les équations différentielles dont l'intégrale générale est uniforme*, Bull. Société Mathématique de France. **28** (1900), 201–261.

[2] P. Painlevé, *Sur les équations différentielles du second ordre et d'ordre supérieur dont l'intégrale est uniforme*, Acta Math. **25** (1902), 1–85.

[3] B. Gambier, *Sur les équations différentielles du second ordre et du premier degré dont l'intégrale générale est à points critiques fixes*, Acta Math. **33** (1910), 1–55.

[4] C. M. Cosgrove and G. Scoufis, *Painlevé classification of a class of differential equations of the second order and second degree*, Studies in Applied Mathematics. **88** (1993), 25-87.

[5] C. M. Cosgrove, *All binomial-type Painlevé equations of the second order and degree three or higher*, Studies in Applied Mathematics. **90** (1993), 119-187.

[6] F. Bureau, *Integration of some nonlinear systems of ordinary differential equations*, Annali di Matematica. **94** (1972), 345–359.

[7] J. Chazy, *Sur les équations différentielles dont l'intégrale générale est uniforme et admet des singularités essentielles mobiles*, Comptes Rendus de l'Académie des Sciences, Paris. **149** (1909), 563–565.

[8] J. Chazy, *Sur les équations différentielles dont l'intégrale générale posséde une coupure essentielle mobile* , Comptes Rendus de l'Académie des Sciences, Paris. **150** (1910), 456–458.

[9] J. Chazy, *Sur les équations différentielles du trousiéme ordre et d'ordre supérieur dont l'intégrale a ses points critiques fixes*, Acta Math. **34** (1911), 317–385.

[10] Y. Sasano, *Coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of types* $B_6^{(1)},D_6^{(1)}$ *and* $D_7^{(2)}$, preprint.

[11] Y. Sasano, *Four-dimensional Painlevé systems of types* $D_5^{(1)}$ *and* $B_4^{(1)}$, preprint.

[12] Y. Sasano, *Higher order Painlevé equations of type* $D_l{}^{(1)}$, RIMS Kokyuroku **1473** (2006), 143–163.

[13] Y. Sasano, *Symmetries in the system of type* $D_4^{(1)}$, preprint.

[14] Y. Sasano, *Coupled Painlevé III systems with affine Weyl group symmetry of types* $B_4^{(1)}$*,* $D_4^{(1)}$ *and* $D_5^{(2)}$, preprint.

[15] Y. Sasano, *Coupled Painlevé III systems with affine Weyl group symmetry of types* $B_5^{(1)},D_5^{(1)}$ *and* $D_6^{(2)}$, preprint.

[16] Y. Sasano, *Coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of type* $D_6^{(1)}$*, II*, RIMS Kokyuroku Bessatsu. **B5** (2008), 137–152.

[17] Y. Sasano, *Coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of type* $E_6^{(2)}$, preprint.

[18] Y. Sasano, *Symmetry in the Painlevé systems and their extensions to four-dimensional systems*, Funkcial. Ekvac. **51** (2008), 351–369.